

120-300-397

BSFB
Lot No.:
Exp. Date:

**EACH TABLET
CONTAINS:** 25 mg
acepromazine maleate.

INDICATIONS: As an aid in
tranquilization and as a
preanesthetic agent in dogs.

DOSAGE: Dogs - 0.25-1.0 mg
per lb of body weight. Dosage
may be repeated as required.

Refer to folded package
insert attached to this
container for complete
product information.

NDC 57319-446-15

Acepromazine Maleate Tablets, USP

25 mg per Tablet

FOR USE IN DOGS ONLY.

CAUTION:
Federal
law restricts
this drug to
use by or on
the order of a
licensed
veterinarian.

**Net Contents:
500 Tablets**

PHOENIX™
Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507



670230-00
80732279
80728743, R.0

STORAGE: Store at 20° to
25°C (68° to 77°F),
excursions permitted between
15° and 30°C (between 59°
and 86°F).

**Keep out of reach of
children.**

Trademarks are property
of Clipper Distributing
Company, LLC.

NADA 117-532, Approved by FDA
Acepromazine Maleate Tablets, USP


For use in dogs only.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION: Acepromazine maleate, a potent neuroleptic agent with a low order of toxicity, is of particular value in the tranquilization of dogs. Its rapid action and lack of hypnotic effect are added advantages.

CHEMISTRY: 2-acetyl-10-(3-dimethylaminopropyl) phenothiazine hydrogen maleate.

Acepromazine Maleate



MODE OF ACTION: Acepromazine maleate has a depressant effect on the central nervous system and therefore causes sedation, muscular relaxation and a reduction in spontaneous activity. It acts rapidly, exerting a prompt and pronounced calming effect. It is an effective preanesthetic agent and lowers the dosage requirement of general anesthetics.

INDICATIONS: As an aid in tranquilization and as a preanesthetic agent in dogs.

Acepromazine Maleate Tablets can be used as an aid in controlling intractable animals during examination, treatment, grooming, x-ray and minor surgical procedures.

DOSEAGE AND ADMINISTRATION: Dogs: 0.25-1.0 mg/lb of body weight. Dosage may be repeated as required.

ANIMAL SAFETY: Acute and chronic toxicity studies have shown a very low order of toxicity for acepromazine maleate.

A safety study using elevated dosages of acepromazine maleate demonstrated no adverse reactions even when administered at three times the upper limit of the recommended daily dosage (3.0 mg/lb body weight). The clinical observation for this high dosage was mild depression which disappeared in most dogs 24 hours after termination of dosing.

The only occurrence of adverse reaction during numerous clinical trials was a very mild respiratory distress (reverse sneeze) which was transient in nature and had no effect on the desired action of the drug.

CONTRAINDICATIONS: Phenothiazines may potentiate the toxicity of organophosphates. Therefore, do not use acepromazine maleate to control tremors associated with organic phosphate poisoning.

Do not use in conjunction with organophosphorus vermifuges or ectoparasiticides, including flea collars. Do not use with procaine hydrochloride.

WARNING: Do not use in animals intended for human consumption.

PRECAUTIONS: Tranquilizers are potent central nervous system depressants, and they can cause marked sedation with suppression of the sympathetic nervous system. Tranquilizers can produce prolonged depression or motor restlessness when given in excessive amounts or when given to sensitive animals. Tranquilizers are additive in

action to the actions of other depressants and will potentiate general anesthesia. Tranquilizers should be administered in smaller doses and with greater care during general anesthesia and also to animals exhibiting symptoms of stress, debilitation, cardiac disease, sympathetic blockade, hypovolemia or shock. Acepromazine, like other phenothiazine derivatives, is detoxified in the liver; therefore, it should be used with caution on animals with a previous history of liver dysfunction or leukopenia.

Epinephrine is contraindicated for treatment of acute hypotension produced by phenothiazine-derivative tranquilizers since further depression of blood pressure can occur.

Phenothiazines should be used with caution when followed by epidural anesthetic procedures because they may potentiate the arterial hypotensive effects of local anesthetics.

ADVERSE REACTIONS: A few rare but serious occurrences of idiosyncratic reactions to acepromazine may occur in dogs following oral or parenteral administration. These potentially serious adverse reactions include behavioral disorders in dogs such as aggression, biting/chewing, and nervousness.

To report suspected adverse reactions, to obtain a Material Safety Data Sheet or for technical assistance, call 1-866-638-2226.

STORAGE: Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F).

HOW SUPPLIED: Acepromazine Maleate Tablets are available in 10 & 25 mg concentrations, and are quarter scored for convenience of administration. Both concentrations are available in bottles of 100 and 500 tablets.

NDC 57319-445-13 - 10 mg - 100 tablets
NDC 57319-445-15 - 10 mg - 500 tablets
NDC 57319-446-13 - 25 mg - 100 tablets
NDC 57319-446-15 - 25 mg - 500 tablets

670230-00
80732279
80728743, R.0
Revised 04/2012

EACH TABLET CONTAINS: 25 mg acepromazine maleate.

INDICATIONS: As an aid in tranquilization and as a preanesthetic agent in dogs.

DOSEAGE: Dogs - 0.25-1.0 mg per lb of body weight. Dosage may be repeated as required.

Refer to folded package insert attached to this container for complete product information.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

PHOENIX™
Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

Net Contents:
500 Tablets



670230-00
80732279
80728743, R.0

NDC 57319-446-15

Acepromazine Maleate Tablets, USP

25 mg per Tablet

FOR USE IN DOGS ONLY.

STORAGE: Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F).

Keep out of reach of children.

Trademarks are property of Clipper Distributing Company, LLC.

TEAR HERE



3 57319 44615 0